



离散数学 (011122)



魏可佶

kejiwei@tongji.edu.cn
<https://kejiwei.github.io/>



- 4.1 Definition and Representation of Relations
- 4.2 Relational Operations
- 4.3 Properties of Relations
- 4.4 Equivalence Relations and Partial Order Relations

- 4.1.1 Ordered Pairs and Cartesian Product
- 4.1.2 Definition of Binary Relations
- 4.1.3 Representation of Binary Relations

■ Definition 4.1- Ordered Pairs

A pair of elements x and y arranged in a specific order is called an *ordered pair* (or sequence), denoted as $\langle x, y \rangle$

e.g. >>> Example:

The Cartesian coordinates of a point: $(3, -4)$

■ Properties of Ordered Pairs

- Order $\langle x, y \rangle \neq \langle y, x \rangle$ (when $x \neq y$)
- The necessary and sufficient condition for $\langle x, y \rangle = \langle u, v \rangle$ is $\langle x, y \rangle = \langle u, v \rangle \Leftrightarrow x = u \wedge y = v$

e.g. >>> Example: $\langle 2, x+5 \rangle = \langle 3y-4, y \rangle$, 求 x, y .

Solve: $3y-4=2, x+5=y \Rightarrow y=2, x=-3$

■ Definition 4.2- Cartesian product

Let A and B be sets, the Cartesian product of A and B is denoted as $A \times B$,

$$A \times B = \{ \langle x, y \rangle \mid x \in A \wedge y \in B \}.$$

e.g. >>> Example: $A = \{0, 1\}$, $B = \{a, b, c\}$

$$A \times B = \{ \langle 0, a \rangle, \langle 0, b \rangle, \langle 0, c \rangle, \langle 1, a \rangle, \langle 1, b \rangle, \langle 1, c \rangle \}$$

$$B \times A = \{ \langle a, 0 \rangle, \langle b, 0 \rangle, \langle c, 0 \rangle, \langle a, 1 \rangle, \langle b, 1 \rangle, \langle c, 1 \rangle \}$$

- Empty Set Interaction: If either A or B is an empty set, then $A \times B$ is the empty set. $A \times \emptyset = \emptyset \times B = \emptyset$
- Not suitable for the commutative property:
 $A \times B \neq B \times A$ ($A \neq B$, $A \neq \emptyset$, $B \neq \emptyset$)
- Not suitable for the associative property:
 $(A \times B) \times C \neq A \times (B \times C)$ ($A \neq \emptyset$, $B \neq \emptyset$, $C \neq \emptyset$)
- The union and intersection operations satisfy the distributive property.
 $A \times (B \cup C) = (A \times B) \cup (A \times C)$
 $(B \cup C) \times A = (B \times A) \cup (C \times A)$
 $A \times (B \cap C) = (A \times B) \cap (A \times C)$
 $(B \cap C) \times A = (B \times A) \cap (C \times A)$
- Cardinality Multiplication Property:
If $|A| = m$, $|B| = n$, then $|A \times B| = mn$

■ Definition 4.3: n-dimensional Cartesian product

(1) An ordered n-tuple is formed by arranging n elements

x_1, x_2, \dots , in a specific order, denoted as $\langle x_1, x_2, \dots, x_n \rangle$

(2) Let A_1, A_2, \dots , be sets., Then the Cartesian product

$$A_1 \times A_2 \times \dots \times A_n = \{ \langle x_1, x_2, \dots, x_n \rangle \mid x_i \in A_i, i=1, 2, \dots, n \}$$

is called the *n-dimensional* (n-ary) *Cartesian product*.

e.g. >>> Example:

(1, 1, 0) is the Cartesian coordinate of a point in space,

$(1, 1, 0) \in R \times R \times R$ (1, 1, 0).

↳ 4.1.2 Definition of Binary Relations

■ Definition 4.4- Binary Relations

A set is called a *binary relation*, denoted as R , if it satisfies one of the following conditions:

- (1) The set is non-empty, and its elements are ordered pairs.
- (2) The set is empty.
 - For example, if $\langle x, y \rangle \in R$, it can be written as $x R y$; if $\langle x, y \rangle \notin R$, it can be written as $x \not R y$

e.g. >>> Example:

$$R = \{\langle 1, 2 \rangle, \langle a, b \rangle\}, S = \{\langle 1, 2 \rangle, a, b\}$$

- R is a binary relation, but S is not a binary relation because a and b are not ordered pairs.
- Using the above notation, we can write $1R2$, aRb , $a \not R c$, etc.

↳ 4.1.2 Definition of Binary Relations (e.g.)

e.g. >>> Example:

$$(1) R = \{ \langle x, y \rangle \mid x, y \in \mathbb{N}, x + y < 3 \}$$
$$= \{ \langle 0, 0 \rangle, \langle 0, 1 \rangle, \langle 0, 2 \rangle, \langle 1, 0 \rangle, \langle 1, 1 \rangle, \langle 2, 0 \rangle \}$$

(2) $C = \{ \langle x, y \rangle \mid x, y \in \mathbb{R}, x^2 + y^2 = 1 \}$, Where R represents the set of real numbers, C is the relation between the horizontal and vertical coordinates of points in the Cartesian coordinate plane, and all points in C exactly form the unit circle in the coordinate plane.

$$(3) R = \{ \langle x, y, z \rangle \mid x, y, z \in \mathbb{R}, x + 2y + z = 3 \},$$

R represents a plane in the 3D Cartesian coordinate system.

↳ 4.1.2 Definition of Binary Relations (e.g.)

The employee payroll (relation) is a set of *5-tuples representing* employees: $\langle 301, B, 25, M, 19000 \rangle$, $\langle 302, O, 23, F, 18500 \rangle$

ID No.	Name	Age	Gender	Salary
301	B	25	M	19000
302	O	23	F	18500
303	Y	37	M	25000
304	M	31	M	22000
...

↳ Binary relation from A to B & Binary relation on A

■ Definition 4.5

Let A and B be sets. Any subset of $A \times B$ that defines a binary relation is called a *binary relation from A to B*. When $A=B$, it is called a *binary relation on A*.

■ Counting:

- $|A|=n, |B|=m, |A \times B|=nm$, and there are 2^{nm} subsets of $A \times B$. Therefore, there are 2^{nm} different binary relations from A to B .
- If $|A|=n$, there are 2^{n^2} different binary relations on A .
- For example, if $|A|=3$, then there are 512 different binary relations on A .

📄 **Note:** Sets are the static carriers of relations, while relations are the dynamic interaction rules between sets.

↳ universal relation(E_A) & identity relation (I_A)

- Let A be any set. The empty set \emptyset is considered as the *empty relation* on A .
- Definition 4.6:
 - E_A is called the *universal relation* on A , where
$$E_A = \{ \langle x, y \rangle \mid x \in A \wedge y \in A \} = A \times A$$
 - I_A is called the *identity relation* on A , where
$$I_A = \{ \langle x, x \rangle \mid x \in A \}$$

e.g. >>> Example: $A = \{1, 2\}$, then

$$E_A = \{ \langle 1, 1 \rangle, \langle 1, 2 \rangle, \langle 2, 1 \rangle, \langle 2, 2 \rangle \}$$

$$I_A = \{ \langle 1, 1 \rangle, \langle 2, 2 \rangle \}$$

■ Definition 4.7:

The less than or equal to relation L_A , the divides relation D_B , and the containment relation R_{\subseteq} are defined as follows:

- $L_A = \{ \langle x, y \rangle \mid x, y \in A \wedge x \leq y \}$, $A \subseteq \mathbb{R}$, \mathbb{R} is the set of real numbers.
- $D_B = \{ \langle x, y \rangle \mid x, y \in B \wedge x \text{ divides } y \}$, $B \subseteq \mathbb{Z}^*$, \mathbb{Z}^* is the set of non-zero integers.
- $R_{\subseteq} = \{ \langle x, y \rangle \mid x, y \in A \wedge x \subseteq y \}$, where A is a family of sets..

e.g. >>> Example: $A = \{1, 2, 3\}$, $B = \{a, b\}$, then

$$L_A = \{ \langle 1, 1 \rangle, \langle 1, 2 \rangle, \langle 1, 3 \rangle, \langle 2, 2 \rangle, \langle 2, 3 \rangle, \langle 3, 3 \rangle \}$$

$$D_A = \{ \langle 1, 1 \rangle, \langle 1, 2 \rangle, \langle 1, 3 \rangle, \langle 2, 2 \rangle, \langle 3, 3 \rangle \}$$

4.1.2 Definition of Binary Relations • Less than or equal to relation(L_A) , Divides relation(D_B), Containment relation(R_{\subseteq})

e.g. \ggg Example: $B=\{a,b\}$, $A=P(B)=\{\emptyset,\{a\},\{b\},\{a,b\}\}$,

then the *inclusion relation on A* is :

$$R_{\subseteq}=\{\langle\emptyset,\emptyset\rangle,\langle\emptyset,\{a\}\rangle,\langle\emptyset,\{b\}\rangle,\langle\emptyset,\{a,b\}\rangle,\langle\{a\},\{a\}\rangle,\langle\{a\},\{a,b\}\rangle,\langle\{b\},\{b\}\rangle,\langle\{b\},\{a,b\}\rangle,\langle\{a,b\},\{a,b\}\rangle\}$$

Similarly, greater than or equal to relations, less than relations, greater than relations, proper inclusion relations, etc., can also be defined.

↳ Relational Matrix & Relational Graph

■ Ways to Represent Relations:

Set-theoretic expression of relations; Relational matrix; Relational graph

■ Definition 4.8: Relational Matrix

If $A = \{x_1, x_2, \dots, x_m\}$ and $B = \{y_1, y_2, \dots, y_n\}$ and R is a relation from A to B , then the *relational matrix* of R is the Boolean matrix $M_R = [r_{ij}]_{m \times n}$, where

$$r_{ij} = 1 \Leftrightarrow \langle x_i, y_j \rangle \in R.$$

■ Definition 4.9: Relational Graph

If $A = \{x_1, x_2, \dots, x_m\}$, and R is a relation on A , *the relational graph* of R is $G_R = \langle A, R \rangle$, where A is the set of vertices and R is the set of edges. If $\langle x_i, x_j \rangle$ belongs to R , there is a directed edge from x_i to x_j in the graph.


↳ Relational Matrix & Relational Graph

 **Note:** Let A and B be finite sets.

- **Relational matrices** are suitable for representing relations from A to B or relations on A .
- **Relational graphs** are suitable for representing relations on A .

e.g. **Example:** $A = \{a, b, c, d\}$, $R = \{ \langle a, a \rangle, \langle a, b \rangle, \langle a, c \rangle, \langle b, a \rangle, \langle d, b \rangle \}$,
The relationship matrix M_R and the relationship graph G_R

$$M_R = \begin{bmatrix} 1 & 1 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$



Objective :

Key Concepts :